

THIAMETHOXAM 25% WG

INSECTICIDE



**BROAD-SPECTRUM
CONTROL**



RAPID TRANSLOCATION



**CONTACT & SYSTEMIC
ACTION**



**EXTENDED RESIDUAL
EFFECT**

TARGET PESTS

Neonicotinoid insecticide with systemic, contact, and stomach action providing rapid uptake and translocation throughout plant with extended residual control of sucking pests.

CROPS

- Rice
- Cotton
- Vegetables, Fruits
- Wheat, Sugarcane



Rice, Cotton, Vegetables, Fruits, Wheat, Sugarcane

TARGET PESTS

- Aphis (Aphis spp.)
- Whitefly (*Bemisia tabaci*)
- Jassids (*Amrasca biguttula*)
- Brown plant hopper (*Nilaparvatta lugens*)
- Thrips (Thrips tabaci)

DOSAGE & RECOMMENDATION

CROPS	COMMON DISEASE	DOSAGE /Aa 1.g/ml% _{yr}	DILUTION	WAITING PERIOD TO HARVEST (in days)
			LTR./mL	
Rice (Paddy)	Ster=borrer Effareerrents Green hopper	100	500-750 L	14
Cotton	Whitefly	1000	500-1000 L	21
Chiti	Whitefly, Aphids	500	300 L water	21
Okra	Whitefly S.Jassids	500	300 L water	3

APPLICATION BENEFITS

- Rapid translocation through the plant.
- Extended residual control.
- Systemic and contact action control sucking pests.
- New growth protection.



**STRONG
MANUFACTURING**



**INNOVATIVE
R&D**



**GLOBAL
REGULATORY
SUPPORT**



**DECADES OF
EXPERIENCE**