

# FLOCOUMAFEN 0.05% BLOCK BAIT

## RODENTICIDE



**SECOND-GENERATION**



**SINGLE-FEED  
LETHAL DOSE**



**VITAMIN KINHIBITION**



**STRONG EFFECTIVE  
CONTROL**

### PRODUCT DESCRIPTION

Second-generation anticoagulant rodenticide in palatable blocformulation.

Provides effective control of rodents including those resistant to first-generation anticoagulants with Single-feed efficacy.

### APPLICATION AREAS:

- Agricultural fields, Warehouses, Residential areas,
- Commercial premises

### MODE OF ACTION



**Norway Rat**



**Roof Rat**



**House mouse**




**Bandicoot Rat**


### DOSAGE & RECOMMENDATION

Usage	Common Pest	Dosage/ha a.i (mg)	Formulation in (ml)	PHI
For rodent control in field storage and crops like rice, soybean	<i>Norway rat,</i> <i>Roof rat,</i> <i>House mouse,</i> <i>Bandicota rat</i>	0.75-1.0	At an interval of 5-10m in bait station or active burrow. Repeat the application after 14 days if problem persists.	NA



 **STRONG  
MAND-ACTURING**

 **INNOVATIVE  
R&D**

 **GLOBAL  
REGULATORY  
SUPPORT**